



Package Leaflet for:

Dr. PONSOLD
Pharma GmbH
Tiergesundheit

Ponsocol®

10⁸ CFU/ml Suspension

For animals, species: **Bos taurus (calf)**

Manufacturer's Name and Address

Dr. Ponsold Pharma GmbH, Loerfeldstraße 20, 58313 Herdecke, Germany
Phone: +49 (0)2330 977 677, Fax: (+49) (0) 2330 977 697

Name of the Veterinary Medicinal Product

Ponsocol® 10⁸ CFU/ml Suspension
For animals, species: **Bos taurus (calf)**
Active substance: **E. coli strain Nissle 1917**

Statement of the active Substance and other Ingredients

Active Ingredients

The active substance is: **Escherichia coli strain Nissle 1917**
1 ml suspension contains bacterial culture of **Escherichia coli strain Nissle 1917** corresponding to 10⁸ viable cells (CFU).

Excipients

Purified water, sodium chloride, potassium chloride, magnesium-sulfate, calcium chloride, magnesium-chloride.

Indication

For the prevention (prophylaxis) of diarrhea in newborn calves.

Contraindications

Not known

Adverse reactions


Not known.
If you notice any serious effects or other effects not mentioned in this package leaflet, please inform your veterinary surgeon.


Target species

Bos taurus (calf)

Dosage, Route and Method of Administration

The following instructions are applied if your veterinarian has not made different statements to the administration of Ponsocol. Please, always take Ponsocol® exactly as your veterinarian has told you, because otherwise Ponsocol® cannot be work properly.





After shaking the container 15 ml of Ponsocol® are directly administered into the mouth from the 1st to the 10th day of life (included). On the 1st day of life it is recommended to apply 15 ml of Ponsocol® before the first feeding.

From the 2nd to the 10th day of life 15 ml Ponsocol® should be administered in the morning before feeding.

Advice on correct administration

If you gave more Ponsocol® than you should

Even after taking too large amounts no special measures are necessary.

If you gave too little or forgot to give Ponsocol®

In cases of a too little or forgotten dose of Ponsocol®, the application should be continued without considering the failure of administration, in accordance with the conditions of use and/or the instructions given by your veterinarian.

Waiting Period

Edible calf tissues: 0 days

Special Storage Precautions

Keep out of the sight and reach of children.

Do not use this veterinary medicinal product after the expiry date which is stated on the carton and on the ampoule.

Store between 2 °C and 8 °C! Do not freeze!

Special warnings

Interactions With Other Drugs

Antibiotics against gram-negative bacteria and sulphonamides may reduce the efficacy of Ponsocol®.

Special Precautions for the Disposal of Unused Medicines or Other Special Precautions in Order to Prevent Pollution of the Environment

Unused animal medicines are best collected at central collecting points for pollutants. If disposing such material together with regular household rubbish, any misuse of the substances by third parties must be prevented. Animal medicines must not be disposed together with used water or into the sewers.

Date on which the package leaflet was last approved

October 2015

Other Information

Pack sizes Packs with 1 x 10 x 15 ml
 Packs with 10 x 10 x 15 ml
 Packs with 50 x 10 x 15 ml

For any information about this veterinary medicinal product, please contact the marketing authorisation holder:

Dr. Ponsold Pharma GmbH, Loerfeldstraße 20, 58313 Herdecke, Germany
Phone: +49 (0)2330 977 677, Fax: (+49) (0) 2330 977 697

Dear User!

It is possible that the leaflet in your medicine pack may differ from this version. This leaflet is an internal, unofficial translation of the German package leaflet and may not apply to other countries.

